

Realize optimal yield and quality in brassica/cole crops with reliable and consistent control

DuPont™
Coragen®
 insect control
 powered by
 RYNAXYPYR®

Brassica/Cole Crops — Southeastern U.S.

DuPont™ Coragen® insect control is a breakthrough mode of action insecticide that delivers remarkable protection. Optimize yield and quality in your fields by achieving consistent and long-lasting control of key insect pests in brassica/cole crops.

Key benefits of Coragen®:

- Helps break the insect resistance cycle with a breakthrough mode of action, compared to other standard treatments
- Excellent control of all major worm pests and leafminer (larvae)¹ plus suppression of silverleaf whitefly nymphs
- The first product to include the only active ingredient that controls worms through drip chemigation²
- Long-lasting protection — 14 to 21 days and longer³
- Shortest reentry — 4 hours
- Short preharvest interval — 3 day for brassica/cole crops
- Excellent crop protection — insects stop feeding in minutes
- Application flexibility — drip² or foliar spray (ground and air)
- Works through ingestion or contact to deliver ovicidal⁴ and ovi-larvicidal activity
- Minimal impact on beneficials when applied at labeled rates — excellent fit with IPM programs*
- Does not flare secondary pests or mites
- Rainfast once spray has dried

* In line with Integrated Pest Management and Good Agricultural Practices, insecticide applications should be made when pollinators are not foraging to avoid unnecessary exposure.

¹ Control of *Liriomyza* species, except suppression only for *L. huidabrensis* and *L. langei*.

² Drip chemigation is not allowed on all crops. See Coragen® product label for which crops drip chemigation is allowed.

³ Depending on application method, untreated plant material may not be fully protected as a result of plant growth.

⁴ Ovicidal activity on selected pest species.

⁵ Additional labeled crops include artichoke, asparagus, bulb vegetables, cereal grains, corn (sweet, seed, field, pop), grass forage, fodder and hay, herbs, hops, legume vegetables, peppermint/ spearmint, non-grass animal feeds (including alfalfa), okra, potato, oilseeds, soybeans, spices, strawberry, sugarcane, tobacco and root and tuber vegetables.

Crops ⁵	Pests (partial list)
	Beet armyworm (<i>Spodoptera exigua</i>)
	Cabbage looper (<i>Trichoplusia ni</i>)
	Corn earworm (<i>Helicoverpa zea</i>)
	Leafminers (larvae) (<i>Liriomyza</i> spp.) ¹
	Silverleaf whitefly nymph suppression
	Beet armyworm (<i>Spodoptera exigua</i>)
	European corn borer (<i>Ostrinia nubilalis</i>)
	Leafminers (larvae) (<i>Liriomyza</i> spp.) ¹
	Loopers
	Tomato fruitworm (<i>Helicoverpa zea</i>)
	Tomato pinworm (<i>Keiferia lycopersicella</i>)
	Western yellowstriped armyworm (<i>Spodoptera praefica</i>)
	Silverleaf whitefly nymph suppression
	Beet armyworm (<i>Spodoptera exigua</i>)
	Cabbage looper (<i>Trichoplusia ni</i>)
	Cross-striped cabbageworm (<i>Evergestis rimosalis</i>)
	Diamondback moth (<i>Plutella xylostella</i>)
	Imported cabbageworm (<i>Pieris rapae</i>)
	Beet armyworm (<i>Spodoptera exigua</i>)
	Cabbage looper (<i>Trichoplusia ni</i>)
	Leafminers (larvae) (<i>Liriomyza</i> spp.) ¹
	Melonworm (<i>Diaphania hyalinata</i>)
	Silverleaf whitefly nymph suppression

DuPont™ Coragen®

insect control

powered by
RYNAXYPYR®

Brassica/Cole Crops — Insecticide Spray Program — Southeastern U.S.

Week	Treatment	Insects**
1–2	DuPont™ Avaunt® — 3.5 oz/A foliar spray, application timing based on local thresholds ⁶	Worms, including diamondback moth
2–3	DuPont™ Coragen® — 3.5–5 fl oz/A foliar spray, application timing based on local thresholds	Worms
3–6	Avaunt® — 3.5 oz/A foliar spray, application timing based on local thresholds ⁶	Worms, including cabbage looper, diamondback moth
6–7	Coragen® — 3.5–5 fl oz/A foliar spray, application timing based on local thresholds	Worms
7–8	Coragen® — 3.5–5 fl oz/A foliar spray ⁷ 10–14 days following previous Coragen® application	Worms
8–12	Avaunt® — 3.5 oz/A foliar spray as needed ⁶	Worms, including cabbage looper, diamondback moth, imported cabbageworm
	<i>or</i> DuPont™ Lannate® SP — 0.5–1.0 lb/A foliar spray as needed	Worms, including cabbage looper, ⁸ diamondback moth, imported cabbage worm

** See individual product labels for specific crop/pest combinations controlled or suppressed by these products.

⁶ Add wetting agent to improve coverage.

⁷ See Coragen® product label for specific guidance regarding Insect Resistance Management practices; avoid using the same mode of action on consecutive generations of insect pests.

⁸ Do not use Lannate® SP for loopers in Alabama and Georgia.

For more information

Put this powerful tool to work for you. Contact your local DuPont retailer or representative to learn how you can get more consistent control of key pests in brassica/cole crops with DuPont™ Coragen® insect control. And visit us at coragen.dupont.com.

DuPont™ Lannate® SP is a restricted-use pesticide.

This reference guide is not intended as a substitute for the product label for the product(s) referenced herein. Product labels for the above product(s) contain important precautions, directions for use, and product warranty and liability limitations, which must be read before using the product(s). Applicators must be in possession of the product label(s) at the time of application. Always read and follow all label directions and precautions for use when using any pesticide alone or in tank-mix combinations.

The DuPont Oval Logo, DuPont™, Avaunt®, Coragen®, Lannate® and Rynaxypyr® are trademarks or registered trademarks of DuPont or its affiliates.

Copyright © 2008-2014 E.I. du Pont de Nemours and Company. All Rights Reserved. 10/14

Reorder No.: K-28162 (Replaces K-23036)