

Realize a simple and reliable solution for weed control in grapes

**DuPont™
Matrix® SG**
herbicide

Grapes — Pacific Northwest

Every season, grape growers and crop consultants have to decide on the most-effective weed-control strategies and products for their vineyards and growing areas.

Fortunately, DuPont™ Matrix® SG herbicide provides a simple and reliable solution. Matrix® SG makes the best foundational choice for grape weed control for multiple reasons: commercially proven crop tolerance; PRE and early POST activity; a labeled low-use rate; an easy-to-mix, -measure and -use formulation, broad vine crop labeling and no soil or use period restrictions.

Building your foundational weed-control program on the power of Matrix® SG is the most-effective way to reduce contact-spray applications and avoid residual herbicides that have limited crop labeling, precautionary crop safety concerns, restrictive nonbearing labeling, or application timing and soil restrictions.

Key benefits of Matrix® SG

- Broad crop labeling (grapes, tree nuts, stone fruits, pome fruits, citrus, tomatoes, potatoes, field corn).
- PRE and early POST activity.
- Excellent broadleaf weed and grass control.
- Yellow nutsedge suppression.
- Commercially proven crop tolerance on young trees (one growing season) and long residual weed control.
- An additional mode of action to minimize selection pressure on glyphosate-tolerant weed species.
- No application restrictions during dormant or nondormant periods.
- No soil restrictions.

Prescription weed-management strategy

- Position Matrix® SG at 4 oz/A as your foundational preemergence herbicide and add tank-mix to maximize broad-spectrum pre- and postemergence weed control. Matrix® SG at 4 oz/A provides consistent control of

problem weeds such as filaree, marestalk, fleabane, Italian ryegrass¹, panicle willowweed¹ and suppresses yellow nutsedge.

- Where tough winter annual weeds — such as marestalk, filaree or panicle willowweed¹ — are problems, research has shown that late fall to early winter preemergence applications of Matrix® SG at 4 oz/A (used with other herbicides registered for grapes, such as Goal 2XL, Alion SC or Chateau) provide excellent residual control.
- Late winter to early spring applications of Matrix® SG at 4 oz/A (used with other herbicides registered for grapes, such as Alion SC or Prowl H2O with activating rain or irrigation) extend control of summer annual grasses and broadleaf weeds into the summer months and reduce the need for pre-harvest foliar herbicides.
- Where 50% or less of the vineyard floor is treated in a herbicide band, Matrix® SG may be applied twice a year at 4 oz/A.

Your DuPont representative can assist you with selecting the best tank-mix herbicides for improved burndown and residual weed control for your vineyard.

Application information

- If weeds emerge, add a foliar herbicide for postemergence control of existing weeds.
- Performance is best when weeds are small and treatment areas are free of significant debris/leaves.
- Rainfall or sprinkler irrigation soon after application (within 21 days) maximizes Matrix® SG and other residual herbicide activity.
- In vineyards with drip irrigation, a late fall or early winter application after soil temperatures are below 50°F will allow winter precipitation to incorporate Matrix® SG into the soil.

¹ This Matrix® SG recommendation is permitted under FIFRA Section 2(ee) for preemergence control of additional grass and broadleaf weeds in the states of Arizona, California, Oregon and Washington in labeled vine crops and has not been submitted to or approved by the EPA. The 2(ee) expires on 12/31/2014.

DuPont™ Matrix® SG

herbicide

Weeds controlled or suppressed with Matrix® SG – Preemergence

Broadleaf Weeds			
Burclover ¹ Chamomile, false Chickweed, common ¹ Clover, white ¹ Cocklebur* Dandelion, common (seedling) Dandelion, common (established)* Fiddleneck, coast ¹ Filaree, redstem Filaree, whitestem ¹ Fleabane, hairy Groundsel, common Henbit Kochia Lambsquarters, common* Lettuce, prickly ¹ Mallow, common Malva/Cheeseweed ¹ Maretail/horseweed	Medicago polymorpha <i>Matricaria maritima</i> Stellaria media Trifolium repens <i>Xanthium</i> spp. <i>Taraxacum officinale</i> <i>Taraxacum officinale</i> Amsinckia menziesii <i>Erodium cicutarium</i> Erodium moschatum <i>Conyza bonariensis</i> <i>Senecio vulgaris</i> <i>Lamium amplexicaule</i> <i>Kochia scoparia</i> <i>Chenopodium album</i> Lactuca serriola <i>Malva neglecta</i> Malva parviflora <i>Conyza canadensis</i>	Mustard, birdsrape Mustard, black Pigweed, prostrate* Pigweed, redroot Pigweed, smooth Nightshade, black* Nightshade, hairy* Puncturevine Purslane, common Ragweed, common* Rocket, London ¹ Sweetclover, yellow ¹ Sowthistle, annual ¹ Spurge, prostrate Spurge, spotted Swinecress ¹ Velvetleaf* Willowweed, panicle ¹	<i>Brassica rapa</i> <i>Brassica nigra</i> <i>Amaranthus blitoides</i> <i>Amaranthus retroflexus</i> <i>Amaranthus hybridus</i> <i>Solanum nigrum</i> <i>Solanum sarrachoides</i> <i>Tribulus terrestris</i> <i>Portulaca oleracea</i> <i>Ambrosia artemisiifolia</i> Sysymbrium irio Melilotus officinalis Sonchus oleraceus <i>Euphorbia prostrata</i> <i>Euphorbia maculata</i> Coronopus didymus <i>Abutilon theophrasti</i> Epilobium ciliatum
Grasses		Sedges	
Bluegrass, annual ¹ Barnyardgrass Crabgrass, large Foxtail, giant Foxtail, green Foxtail, yellow Quackgrass Ryegrass, Italian ¹ Wheat, volunteer Oat, wild*	Poa annua <i>Echinochloa crus-galli</i> <i>Digitaria sanguinalis</i> <i>Setaria faberi</i> <i>Setaria viridis</i> <i>Setaria glauca</i> <i>Agropyron repens</i> Lolium multiflorum <i>Triticum aestivum</i> <i>Avena fatua</i>	Nutsedge, yellow*	<i>Esculentus, cyperus</i>

* Partial control

¹ This Matrix® SG recommendation is permitted under FIFRA Section 2(ee) for preemergence control of additional grass and broadleaf weeds in the states of Arizona, California, Oregon and Washington in labeled vine crops and has not been submitted to or approved by the EPA. The 2(ee) expires on 12/31/2014.



Trial Location: American Canyon, Calif.
 Conducted by John Roncorni, UC DAVIS Farm Advisor
 Matrix® SG 4.0 oz/A + Chateau at 6 oz/A
 Treatment applied: January 9, 2013
 Photo taken: June 24, 2013 (166 DAT)
 Primary weeds present: Willowherb, filaree and malva



Trial Location: Madera, Calif.
 Conducted by Ron Vargus, UC Davis Emeritus
 Matrix® SG 4.0 oz/A + Alion 3.75 fl oz/A
 Treatment applied: January 18, 2012;
 Photo taken: May 31, 2012 (134 DAT)
 Primary weed present: Hairy fleabane

For more information

Contact your local DuPont retailer, representative or crop consultant to learn more about Matrix® SG herbicide. And visit us at matrix.dupont.com.