

Realize optimal yield and quality in cantaloupes and melons with reliable and consistent control.

DuPont™

powered by RYNAXYPYR

Cantaloupes and Melons — Eastern U.S.

DuPont™ Coragen® insect control is a breakthrough mode of action insecticide that delivers remarkable protection. Optimize yield and quality in your fields by achieving consistent and long-lasting control of key insect pests in cantaloupes and melons.

Key benefits of Coragen®

- Helps break the insect resistance cycle with a breakthrough mode of action, compared to other standard treatments
- Excellent control of all major worm pests and leafminer (larvae)1 plus suppression of silverleaf whitefly nymphs
- The first product to include the only active ingredient that controls worms through drip chemigation²
- Long-lasting protection 14 to 21 days and longer³
- Shortest reentry 4 hours
- Short preharvest interval -1 day for cantaloupes and melons
- Excellent crop protection insects stop feeding in minutes
- Application flexibility drip² or foliar spray (ground and air)
- Works through ingestion or contact to deliver ovicidal⁴ and ovi-larvicidal activity
- Minimal impact on beneficials when applied at labeled rates — excellent fit with IPM programs*
- Does not flare secondary pests or mites
- Rainfast once spray has dried
- In line with Integrated Pest Management and Good Agricultural Practices, insecticide applications should be made when pollinators are not foraging to avoid unnecessary exposure.
- Control of *Liriomyza* species, except suppression only for *L. huidabrensis* and *L. langei*.

 Drip chemigation is not allowed on all crops. See Coragen® product label for which crops drip chemigation is allowed.
- Depending on application method, untreated plant material may not be fully protected as a result of plant growth.
- Ovicidal activity on selected pest species. Additional labéled crops include artichoke, asparagus, bulb vegetables, cereal grains, corn (sweet, seed, field, pop), grass forage, fodder and hay, herbs, hops, legume vegetables, peppermint/ spearmint, non-grass animal feeds (including alfalfa), okra, potato, oilseeds, soybeans, spices, strawberry, sugarcane, tobacco and root and tuber vegetables.

Crops ⁵	Pests (partial list)
Leafy vegetables (non-brassica)	Beet armyworm (Spodoptera exigua)
	Cabbage looper (Trichoplusia ni)
	Corn earworm (Helicoverpa zea)
	Leafminers (larvae) (<i>Liriomyza</i> spp.)¹
	Silverleaf whitefly nymph suppression
Fruiting vegetables	Beet armyworm (Spodoptera exigua)
	European corn borer (Ostrinia nubilalis)
	Leafminers (larvae) (<i>Liriomyza</i> spp.)¹
	Loopers
	Tomato fruitworm (Helicoverpa zea)
	Tomato pinworm (Keiferia lycopersicella)
	Western yellowstriped armyworm (Spodoptera praefica)
	Silverleaf whitefly nymph suppression
Brassica vegetables	Beet armyworm (Spodoptera exigua)
High the state	Cabbage looper (Trichoplusia ni)
	Cross-striped cabbageworm (Evergestis rimosalis)
	Diamondback moth (Plutella xylostella)
	Imported cabbageworm (Pieris rapae)
Cucurbits	Beet armyworm (Spodoptera exigua)
	Cabbage looper (Trichoplusia ni)
	Leafminers (larvae) (<i>Liriomyza</i> spp.)¹
	Melonworm (Diaphania hyalinata)
	Silverleaf whitefly nymph suppression



DuPont™Coragen® POWERED BY R Y N A X Y P Y R®

Cantaloupes and Melons — Insecticide Spray Program — Eastern U.S.		
Week	Treatment	Insects**
1–5 DuPont™ Avaunt® — 3.5 oz/A foliar spray or DuPont™ Lannate® SP — 0.5–1.0 lb/A foliar spray and/or DuPont™ Vydate® L — 1–2 qt/A via drip chemigation		Pickleworm, melonworm
	DuPont™ Lannate® SP — 0.5–1.0 lb/A foliar spray	Worms, melon aphid, cucumber beetle, flea beetle
	Nematodes ⁶	
5-6 DuPont™ Coragen® — 5 fl oz/A via drip chemigation if applicable; otherwise, foliar spray Vydate® L — 1-2 qt/A via drip chemigation or Vydate® L — 1-2 qt/A foliar spray	DuPont™ Coragen® — 5 fl oz/A via drip chemigation if applicable; otherwise, foliar spray	Worms, including pickleworm, melonworm, plus leafminer (larvae), silverleaf whitefly nymphs s
	Vydate $^{\circ}$ L $-$ 1 $-$ 2 qt/A via drip chemigation	Nematodes ⁶
		Aphids, thrips, nematodes ⁶
Coragen® application	Coragen® — 5 fl oz/A foliar spray³ 14 days after previous Coragen® application	Worms, including pickleworm, melonworm, plus leafminer (larvae), silverleaf whitefly nymphs
	Vydate® L — 1–2 qt/A foliar spray	Aphids, thrips, nematodes ⁶
8+	Avaunt® — 3.5 oz/A foliar spray	Pickleworm, melonworm
	Lannate® SP — 0.5–1.0 lb/A foliar spray	Worms, melon aphid, cucumber beetle, flea beetle
		Aphid, thrips, nematodes ⁶

** See individual product labels for specific crop/pest combinations controlled or suppressed by these products.

- NOTE: Studies have shown that, when used in a program for pest management, application of Vydate® L can result in enhanced plant vigor.

 For supplemental treatment of specific nematode species following a pre-plant or an at-plant nematicide application. See Vydate® L label for details.
- Control of Liriomyza species, except suppression only for L. huidabrensis and L. langei.
- Suppression only.
- See Coragen® product label for specific guidance regarding Insect Resistance Management practices; avoid using the same mode of action on consecutive generations of insect pests.

For more information

Put this powerful tool to work for you. Contact your local DuPont retailer or representative to learn how you can get more consistent control of key pests in cantaloupes and melons with DuPont™ Coragen® insect control. And visit us at coragen.dupont.com.