



The miracles of science™

Count on DuPont to help optimize alfalfa yield and pasture quality through reliable and consistent insect control

DuPont™
Prevathon®
insect control
powered by
RYNAXYPYR®

Recommendation Sheet — Alfalfa, Pasture and Grasses

DuPont™ Prevathon® insect control is labeled for use on the full grass and non-grass animal feed crop groups. Help protect the yield and quality of animal feed by achieving reliable and consistent control of worm pests. Prevathon® is highly potent and efficacious and yet at the same time has a favorable mammalian (such as livestock) toxicological profile.

Key Benefits of Prevathon®:

- Delivers long-lasting residual control of the key worm species
- Helps break the insect resistance cycle due to its alternate mode of action versus current standards
- Delivers longer-lasting insect control, which helps eliminate the need for multiple applications per season
- Applications can be made right up to cutting — 0 Day PHI
- Grazing and haying intervals are 0 days — livestock can re-enter and graze immediately after application
- Offers effective control of worm pests while having minimal impact on beneficials such as honeybees†
- Has an excellent worker protection profile, with a 4-hour re-entry period and minimal PPE, which allows you to better manage your crop and your workload

† When used in accordance with label directions.





The miracles of science™

DuPont™ Prevathon® insect control

DuPont™ Prevathon® Use Rates — Alfalfa

Crop	Target Pests	Rate of Application		Last application (Days to Harvest)	REI (Hours)
		Pound active ingredient per acre	Fluid ounces per acre		
Pasture — Crop Group 18 (non-grass animal feeds — forage, fodder, straw and hay)	Alfalfa looper, beet armyworm	0.047 – 0.067	14.0 – 20.0	0	4
Make no more than 4 applications per acre per crop. Make one application per cutting. Do not apply more than 60 fl oz Prevathon® or 0.2 lbs a.i. of chlorantraniliprole-containing products per acre per year.					
Pasture — Crop Group 17 (grass forage, fodder and hay)	Beet armyworm, corn earworm, fall armyworm, southern armyworm	0.047 – 0.067	14.0 – 20.0	0	4
Make no more than 4 applications per acre per crop. Minimum interval between treatments is 7 days. Do not apply more than 60 fl oz Prevathon® or 0.2 lbs a.i. of chlorantraniliprole-containing products per acre per year.					

For more information

Put this powerful tool to work for you. Contact your local crop protection retailer or DuPont representative to learn how you can get more consistent control of key pests with DuPont™ Prevathon® insect control. And visit us on the Web at prevathon.dupont.com.

Prevathon® may not be approved for use on pasture in all states. See your local DuPont retailer or representative for details and availability in your state. This reference guide is not intended as a substitute for the product label for the product(s) referenced herein. Product labels for the above product(s) contain important precautions, directions for use and product warranty and liability limitations that must be read before using the product. Applicators must be in possession of the product label(s) at the time of application. Always read and follow all label directions and precautions for use when using any pesticide alone or in tank-mix combinations. The DuPont Oval Logo, DuPont™, The miracles of science™, Prevathon® and Rynaxypyr® are trademarks or registered trademarks of DuPont or its affiliates. Copyright © 2011-2012 E.I. du Pont de Nemours and Company. All Rights Reserved. 12/12 Reorder No.: K-26235 (Replaces K-24425)