



# DISTILLASE® CS

## Saccharifying Enzyme for Ethanol Production

### DESCRIPTION

DISTILLASE® CS enzyme is an optimized, concentrated blend of enzymes that produces glucose from liquefied starch. DISTILLASE® CS enzyme contains fungal 1,4- $\alpha$ -Dglucan glucohydrolase (E.C. 3.2.1.3) commonly referred to as glucoamylase or amyloglucosidase and fungal alpha-amylase (E.C.3.2.1.1). Both enzymes are derived from recombinant strains of *Trichoderma reesei*. The glucoamylase in DISTILLASE® CS enzyme catalyzes the release of successive glucose units from the nonreducing ends of soluble dextrans and oligosaccharides, while the alpha-amylase at the same time can generate those oligosaccharides and dextrans.

### TYPICAL CHARACTERISTICS

**Activity:** 570 GAU/g (minimum)

**Appearance:** Clear brown liquid

**pH:** 4.0 - 4.5

**Specific gravity:** 1.13 - 1.16 g/ml

**Formulation:** Food grade

### UNIT DEFINITION

One Glucoamylase Unit (GAU) is the amount of enzyme that will liberate one gram of reducing sugars calculated as glucose per hour from soluble starch substrate under the conditions of the assay. A detailed assay method is available upon request.

### PERFORMANCE BENEFITS

DISTILLASE® CS enzyme provides the following benefits to ethanol producers:

- Increased ethanol yield by complete saccharification of liquefied starch
- Less unfermented starch in stillage and DDGS.
- Improved performance in pre-saccharification processes
- Increased fermentation rates and shorter fermentation times

### APPLICATION RECOMMENDATIONS


DISTILLASE® CS enzyme is used to saccharify liquefied starch from various sources including wheat, corn, sorghum, barley, rye, rice and cassava. The resultant glucose product is fermented by yeast to yield ethanol.

#### RECOMMENDED OPERATIONAL CONDITIONS


Optimum pH range	4.0 - 5.0
Effective pH range	3.2 - 5.8
Effective temperature range	28 - 65°C (83 - 149°F)
Pre-saccharification	1 - 16 hours at 60 - 65°C (140 - 149°F)
DISTILLASE® CS dosage	Please contact your DuPont Technical Representative for recommendations specific to your liquefaction process

### DOSAGE GUIDELINES

DISTILLASE® CS enzyme is generally added at a level of 0.055 to 0.075 % w/w (starch, dry solid basis). This corresponds to a level of 0.375 to 0.50 kg DISTILLASE® CS / MT grain 'as is'. The actual enzyme requirement is dependant on the temperature, reaction time and pH of the individual plant saccharification/fermentation process, and its desired results. Fermentation performance may be enhanced by the addition of an acid fungal protease like FERMGEN™.



Effect of initial fermentation pH on conversion efficiency with DISTILLASE® CS enzyme (whole ground wheat, 30% DS, 32°C)



Effect of DISTILLASE® CS enzyme and conventional glucoamylase on residual starch of fermented wheat mash

## COMPLETE SACCHARIFICATION OF STARCH

DISTILLASE® CS enzyme converts more starch to fermentable sugars compared to other comparable products.

## PACKAGING

DISTILLASE® CS enzyme is available in various packaging sizes. Please contact DuPont for detailed information.

## REGULATORY STATUS

This product complies with the current recommended purity specifications for food-grade enzymes given by the Joint FAO/WHO Expert Committee on Food Additives (JECFA) and the Food Chemicals Codex (FCC) and is GRAS (Generally Recognized As Safe) in the United States for use in carbohydrate processing.

## STORAGE

To ensure maximum retention of activity, store DISTILLASE® CS enzyme under refrigerated conditions with the container closed. Prolonged storage at elevated temperature should be avoided.

## SAFETY & ENZYME HANDLING

Inhalation of enzyme dust and mists should be avoided. In case of contact with the skin or eyes, promptly rinse with water for at least 15 minutes.

For detailed handling information, please refer to the appropriate Material Safety Data Sheet, the Enzyme Technical Association (ETA) handbook *Working Safely With Enzymes*, and the Association of Manufacturers and Formulators of Enzyme Products (AMFEP) handbook *Guide to the Safe Handling of Microbial Enzyme Preparations*. All are available from DuPont.

## TECHNICAL SERVICE

Information covering specific applications of this product is available. DuPont will work with customers to enhance processes and solve problems. Let us know what you need and we will assist you.

## CONTACT INFORMATION

### NORTH AMERICA

Rochester, New York (USA)  
 ☎ +1 800 847 5311  
 ☎ +1 585 256 5295

### ASIA/PACIFIC

Singapore  
 ☎ +65 6511 5600  
 ☎ +65 6511 5666

### EUROPE, MIDDLE EAST & AFRICA

Leiden, The Netherlands  
 ☎ +31 71 5686 168  
 ☎ +31 71 5686 169

Shanghai, P.R. China

☎ +86 21 2307 9588  
 ☎ +86 21 2307 9599

### LATIN AMERICA

Buenos Aires, Argentina  
 ☎ +54 11 4875 9500  
 ☎ +54 11 4875 9529

Mumbai, India

☎ +91 22 3008 7131  
 ☎ +91 22 3008 7150

Lahore, Pakistan

☎ +92 300 8476 404  
 ☎ +92 423 5437 866

### SÃO PAULO, BRAZIL

☎ +55 11 4613 3800  
 ☎ +55 11 4612 1101

[www.dupont.com](http://www.dupont.com)

Copyright © 2012 DuPont. All rights reserved. The DuPont Oval Logo, DuPont, the Leaf Globe, and DISTILLASE and FERMGEN are trademarks or registered trademarks of E. I. du Pont de Nemours and Company or its affiliates.

USE AND/OR SALE OF THIS PRODUCT MAY BE COVERED BY ONE OR MORE PATENTS IN THE U.S. AND OTHER COUNTRIES, AS WELL AS PENDING PATENT APPLICATIONS IN THE U.S. AND OTHER COUNTRIES.

The information contained in this product literature is, to the best of our knowledge, true and accurate and the product as sold is in conformance with the specifications set forth herein as determined by the assay methods described herein. Due to conditions of use, technical errors or omissions, improper handling or storage beyond our control, Danisco US Inc. hereby DISCLAIMS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING THE IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Danisco US Inc. shall not be liable for any incidental, consequential, or special damages resulting in any way from the furnishing, performance, or use of this product literature or the product described herein.

Nothing contained herein shall be construed as a representation by Danisco US Inc. that the use or resale of the product or processes described herein will not violate any rules or regulations of any countries, regions, localities, etc. or infringe upon patents or other intellectual property rights of third parties or that the recommendations and usage suggestions described constitute any authorization or inducement to infringe any such rights.

This document is subject to change without further notice.

