

# FERMGEN™ 2.5X

Acid Fungal Protease Enzyme for Dry Grind Ethanol Production

## DESCRIPTION

FERMGEN™ 2.5X is an acid proteolytic enzyme characterized by its ability to hydrolyze proteins under low pH conditions. The broad substrate specificity of FERMGEN™ 2.5X protease enables the enzyme to efficiently hydrolyze most grain proteins in a random fashion.

## TYPICAL CHARACTERISTICS

**Activity:** 2500 SAPU/g (minimum)

**Appearance:** Brown liquid

**pH:** 4.0 - 5.0

**Specific gravity:** 1.12 - 1.20 g/ml


The activity of FERMGEN™ 2.5X protease is expressed in Spectrophotometric Acid Protease Units (SAPU). One SAPU is the amount of enzyme activity that liberates one micromole of tyrosine per minute from a casein substrate under conditions of the assay. A detailed assay method is available upon request.

## PERFORMANCE BENEFITS

FERMGEN™ 2.5X acid fungal protease provides the following benefits to ethanol producers:


- Higher yields and faster fermentation rates for corn, milo and wheat mashes. This is the result of starch freed from protein matrices and enhanced yeast nutrition by specific amino acids as well as di- and tripeptides.
- Reduces or eliminates the need for emulsion breakers in back end corn oil recovery systems.
- Reduces the amount of Nitrogen (Urea or Anhydrous Ammonia) needed in fermentation.
- Enhances yeast propagation and yeast performance.

## EFFECT OF FERMGEN™ 2.5X ON RATE AND TITER




## EFFECT ON YEAST PROPAGATION

FERMGEN™ 2.5X significantly increases the rate of yeast propagation.



Control




FERMGEN™ 2.5X  
Dosage: 0.12 SAPU/g DS

## RECOMMENDED CONDITIONS


FERMGEN™ 2.5X protease functions effectively during typical simultaneous saccharification and fermentation (SSF) conditions: pH 3.0 to 5.0 and temperatures as low as 28°C. The product should be added in a single dose as early as possible to the propagation tank or the fermentor.

RECOMMENDED OPERATIONAL CONDITIONS	
pH	3.5 to 5.0
Dry Substance	26 - 38%
Temperature	The enzyme is effective up to 65°C, but works well at fermentation temperatures of 28 - 35°C

FERMGEN™ 2.5X activity pH profile



FERMGEN™ 2.5X temperature activity profile



## DOSAGE GUIDELINES

The optimal dosage level of FERMGEN™ 2.5X protease in simultaneous saccharification and fermentation is dependent upon processing parameters and the benefit being sought.

DOSE WT/WT AS IS GRAIN	BENEFITS
0.0006% - 0.0012%	Yield Enhancement
0.0017% - 0.0020%	Yield Enhancement Oil Emulsion Reduction
0.0030% - 0.0050%	Yield Enhancement Oil Emulsion Reduction Urea Replacement (75%)*

\*Reductions in urea of up to 75% have been observed in the field. An appropriate starting dosage can be established using your materials at the DuPont Applied Innovation Center.

## PACKAGING

FERMGEN™ 2.5X protease is available in various package sizes. Please consult your DuPont representative for detailed information.

## STORAGE

To ensure maximum retention of activity, store FERMGEN™ 2.5X protease under refrigerated conditions with the container closed.

## SAFETY & ENZYME HANDLING

Inhalation of enzyme dust and mists should be avoided. In case of contact with the skin or eyes, promptly rinse with water for at least 15 minutes.

For detailed handling information, please refer to the appropriate Material Safety Data Sheet, the Enzyme Technical Association (ETA) handbook *Working Safely With Enzymes*, and the Association of Manufacturers and Formulators of Enzyme Products (AMFEP) handbook *Guide to the Safe Handling of Microbial Enzyme Preparations*. All are available from DuPont.

## TECHNICAL SERVICE

Information covering specific applications of this product is available. DuPont will work with customers to enhance processes and solve problems.

## CONTACT INFORMATION

### NORTH AMERICA

Rochester, New York (USA)

☎ +1 800 847 5311

☎ +1 585 256 5295

### ASIA/PACIFIC

Singapore

☎ +65 6511 5600

☎ +65 6511 5666

### EUROPE, MIDDLE EAST & AFRICA

Leiden, The Netherlands

☎ +31 71 5686 168

☎ +31 71 5686 169

Shanghai, P.R. China

☎ +86 21 2307 9588

☎ +86 21 2307 9599

### LATIN AMERICA

Buenos Aires, Argentina

☎ +54 11 4875 9500

☎ +54 11 4875 9529

Mumbai, India

☎ +91 22 3008 7131

☎ +91 22 3008 7150

### SÃO PAULO, BRAZIL

☎ +55 11 4613 3800

☎ +55 11 4612 1101

Lahore, Pakistan

☎ +92 300 8476 404

☎ +92 423 5437 866

[www.dupont.com](http://www.dupont.com)

Copyright © 2013 DuPont. All rights reserved. The DuPont Oval Logo, DuPont, the Leaf Globe, FERMGEN™ and are trademarks or registered trademarks of E. I. du Pont de Nemours and Company or its affiliates.

USE AND/OR SALE OF THIS PRODUCT MAY BE COVERED BY ONE OR MORE PATENTS IN THE U.S. AND OTHER COUNTRIES, AS WELL AS PENDING PATENT APPLICATIONS IN THE U.S. AND OTHER COUNTRIES.

The information contained in this product literature is, to the best of our knowledge, true and accurate and the product as sold is in conformance with the specifications set forth herein as determined by the assay methods described herein. Due to conditions of use, technical errors or omissions, improper handling or storage beyond our control, DuPont and its affiliates hereby DISCLAIMS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING THE IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. DuPont and its affiliates shall not be liable for any incidental, consequential, or special damages resulting in any way from the furnishing, performance, or use of this product literature or the product described herein.

Nothing contained herein shall be construed as a representation by DuPont and its affiliates that the use or resale of the product or processes described herein will not violate any rules or regulations of any countries, regions, localities, etc. or infringe upon patents or other intellectual property rights of third parties or that the recommendations and usage suggestions described constitute any authorization or inducement to infringe any such rights.

This document is subject to change without further notice.

