



# STARGEN™ 002

## Granular Starch Hydrolyzing Enzyme for Ethanol Production

### DESCRIPTION

STARGEN™ 002 enzyme contains *Aspergillus kawachi* alpha-amylase expressed in *Trichoderma reesei* and a gluco-amylase from *Trichoderma reesei* that work synergistically to hydrolyze granular starch substrate to glucose. The endo-activity, alphaamylase and exo-activity, gluco-amylase catalyze the complete hydrolysis of granular starch under a variety of alcohol fermentation conditions.

### TYPICAL CHARACTERISTICS

**Activity:** 570 GAU/g (minimum)

**Appearance:** Clear brown liquid

**Specific gravity:** 1.13 - 1.16 g/ml

**pH:** 4.0 - 4.5

### UNIT DEFINITION

One Glucoamylase Unit (GAU) is the amount of enzyme that will liberate one gram of reducing sugars calculated as glucose per hour from soluble starch substrate under the conditions of the assay. A detailed assay method is available upon request.

### PERFORMANCE BENEFITS

STARGEN™ 002 enzyme provides the following benefits to ethanol producers:

- Hydrolyzes granular starch without the need for starch gelatinization or liquefaction.
- Maintains high activity under simultaneous saccharification and fermentation conditions (pH and temperature) favoured by yeast for ethanol production.

When used in a well designed no-cook ethanol production process and compared to conventional liquefaction and saccharification enzyme system with a cook step it offers:

- Higher ethanol yield per unit of grain
- Reduced energy input per unit of ethanol produced
- Reduced unit operations (e.g. cook, liquefaction)
- Improved sustainability of the production process

### APPLICATIONS

STARGEN™ 002 enzyme is used to saccharify un-cooked grain mashes from various sources including wheat, triticale, rye, barley and rice.

The resultant glucose is fermented by yeast or another appropriate microorganism to yield alcohol. STARGEN™ 002 may be added directly to the fermentor.

STARGEN™ 002 enzyme may be favourably combined with enzyme products like OPTIMASH™ to reduce mash viscosity or FERMGEN™ to provide amino acids and peptides as yeast food.

### RECOMMENDED OPERATIONAL CONDITIONS

pH	3.3-4.5 (To inhibit lactic acid bacteria a low pH might be preferred)
Dry Solids	20-34% w/w
Simultaneous Saccharification and Fermentation	Pretreatment: recommended for all grains to improve hydration and increase conversion rate. The recommended minimum temperature is 48°C (118°F) and is specific for the type of raw material. Fermentation: 24-76 hours at 20-40°C (68-104°F), with sufficient agitation to keep solids suspended.

### DOSAGE GUIDELINES

The optimal usage level of STARGEN™ 002 enzyme in simultaneous saccharification and fermentation is dependant upon processing parameters such as type of raw material, DS (Dry Solids), fermentation time, pH and temperature. A minimum STARGEN™ 002 starting dose of 0.8 - 1.6 kg/MT (metric ton) of grain is recommended under most fermentation conditions. Grains like corn or milo might require a higher dose.

## EFFECT OF GRAIN PARTICLE SIZE

Hammermills are typically used in dry milling of grains. The recommended screen hole diameter are specific for the raw material used.

Screen Hole Diameter	Wheat/Triticale/Rye	Barley	Corn
mm	2.5	3.0	1.5
US mesh	8	6	14

## IMPACT OF SLURRY TEMPERATURE

To improve hydration of milled grains and include hot plant effluents like condensates or thin stillage optimal mixing temperatures for each raw material needs to be adjusted.

Grain	(°C)	(°F)
Rye	49-51	120-124
Wheat	56-57	133-135
Triticale	56-57	133-135
Barley	56-57	133-135
Corn	62-63	144-145


## ENZYMES ADDED TO THE MIXING STEP

To reduce mash viscosity and to activate the starch granules GC 626 (acid alpha amylase) and hemicellulases (OPTIMASH™) should be added to the mixing tank. The recommended pH during mixing is at 3.5-4.0.


Grain	GC 626 Enzyme (kg/MT)	OPTIMASH™ Enzyme (kg/MT)
Rye	0.13	-VR 0.19
Wheat	0.16	-BG 0.10
Triticale	0.13	-VR 0.10
Barley	0.13	-BG 0.20
Corn	0.16	none

## EFFECT OF STARGEN™ 002 ENZYME CONCENTRATION

STARGEN™ 002 enzyme is a highly active product. As the amount of enzyme is increased, the rate of fermentation increases.



Experimental conditions: Substrate: milled whole corn, 28% DS, fermentation temperature: 32°C, 0.20 kg/MT FERMGENT™ protease, 400 ppm urea. Mash pretreated at 60°C for 90 minutes.



Experimental conditions: Substrate: milled whole wheat, 28% DS, fermentation temperature: 32°C, 0.05 kg/MT FERMGENT™ protease, 400 ppm urea. Mash pretreated at 56°C for 90 minutes.

## STORAGE & STABILITY

To ensure maximum retention of activity, store STARGEN™ 002 enzyme under refrigerated conditions with the container closed. Prolonged storage at elevated temperature should be avoided. For more information on the storage of this product, please contact your DuPont representative.

## PACKAGING

STARGEN™ 002 enzyme is available in various packaging sizes. Please contact DuPont for detailed information.

## SAFETY & ENZYME HANDLING

Inhalation of enzyme dust and mists should be avoided. In case of contact with the skin or eyes, promptly rinse with water for at least 15 minutes.

For detailed handling information, please refer to the appropriate Material Safety Data Sheet, the Enzyme Technical Association (ETA) handbook *Working Safely With Enzymes*, and the Association of Manufacturers and Formulators of Enzyme Products (AMFEP) handbook *Guide to the Safe Handling of Microbial Enzyme Preparations*. All are available from DuPont.

## TECHNICAL SERVICE

DuPont will work with customers to enhance processes and solve problems. Let us know what you need and we will assist you.

## CONTACT INFORMATION

### NORTH AMERICA

Rochester, New York (USA)  
 ☎ +1 800 847 5311  
 ☎ +1 585 256 5295

### EUROPE, MIDDLE EAST & AFRICA

Leiden, The Netherlands  
 ☎ +31 71 5686 168  
 ☎ +31 71 5686 169

### LATIN AMERICA

Buenos Aires, Argentina  
 ☎ +54 11 4875 9500  
 ☎ +54 11 4875 9529

### SÃO PAULO, BRAZIL

☎ +55 11 4613 3800  
 ☎ +55 11 4612 1101

### ASIA/PACIFIC

Singapore  
 ☎ +65 6511 5600  
 ☎ +65 6511 5666

### Shanghai, P.R. China

☎ +86 21 2307 9588  
 ☎ +86 21 2307 9599

### Mumbai, India

☎ +91 22 3008 7131  
 ☎ +91 22 3008 7150

### Lahore, Pakistan

☎ +92 300 8476 404  
 ☎ +92 423 5437 866

[www.dupont.com](http://www.dupont.com)

Copyright © 2012 DuPont. All rights reserved. The DuPont Oval Logo, DuPont, the Leaf Globe, STARGEN, FERMGEN are trademarks or registered trademarks of E. I. du Pont de Nemours and Company or its affiliates.

USE AND/OR SALE OF THIS PRODUCT MAY BE COVERED BY ONE OR MORE PATENTS IN THE U.S. AND OTHER COUNTRIES, AS WELL AS PENDING PATENT APPLICATIONS IN THE U.S. AND OTHER COUNTRIES.

The information contained in this product literature is, to the best of our knowledge, true and accurate and the product as sold is in conformance with the specifications set forth herein as determined by the assay methods described herein. Due to conditions of use, technical errors or omissions, improper handling or storage beyond our control, Danisco US Inc. hereby DISCLAIMS ANY EXPRESS OR IMPLIED WARRANTIES, INCLUDING THE IMPLIED WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE. Danisco US Inc. shall not be liable for any incidental, consequential, or special damages resulting in any way from the furnishing, performance, or use of this product literature or the product described herein.

Nothing contained herein shall be construed as a representation by Danisco US Inc. that the use or resale of the product or processes described herein will not violate any rules or regulations of any countries, regions, localities, etc. or infringe upon patents or other intellectual property rights of third parties or that the recommendations and usage suggestions described constitute any authorization or inducement to infringe any such rights.

This document is subject to change without further notice.

