



**SPECIAL LOCAL NEED
FIFRA Section 24(C)LABEL**

**DuPont Crop
Protection**

**DUPONT™CORAGEN®
INSECT CONTROL
HYBRID POPLARS
STATE OF OREGON**

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF OREGON

DUPONT™CORAGEN® INSECT CONTROL

**EPA Reg. No. 352-729
EPA SLN NO. OR-100009**

**FOR USE ON HYBRID POPLARS
FOR CONTROL OF COTTONWOOD LEAF BEETLES**

**THIS LABEL IS VALID UNTIL DECEMBER 31, 2020
OR UNTIL OTHERWISE AMENDED, WITHDRAWN, CANCELED OR SUSPENDED**

Active Ingredient	By Weight
Chlorantraniliprole 3-Bromo-N-[4-chloro-2-methyl-6- [(methylamino)carbonyl]phenyl]-1- (3-chloro-2-pyridinyl)-1H-pyrazole- 5-carboxamide	18.4%
Other Ingredients	81.6%
TOTAL	100.0%

ENVIRONMENTAL HAZARDS

This pesticide is toxic to aquatic invertebrates, oysters, and shrimp. Do not apply directly to water. Drift and runoff may be hazardous to aquatic organisms in water adjacent to use sites. Surface Water Advisory- This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of chlorantraniliprole from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Ground Water Advisory- This chemical has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This Special Local Need (SLN) label and the EPA registered DuPont™ CORAGEN® Insect Control product label must be in the possession of the user at the time of pesticide application.

Follow all applicable directions, restrictions, Worker Protection Standard requirements and precautions on the EPA registered CORAGEN® label.

This use is only for hybrid poplars grown on land classified as cropland by the Farm Service Agency.

USE INFORMATION

DuPont™ CORAGEN® Insect Control is a suspension concentrate that can be applied to control listed insect pests. CORAGEN® is diluted with water for application.

For product information call 1-888-6-DUPONT

© 2016 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Road, Wilmington, DE 19805 U.S.A.

APPLICATION INFORMATION, RATES AND TIMING

CORAGEN® insect control is labeled for the control of Cottonwood Leaf Beetle (*Chrysomela scripta*) adults and larvae when applied as a foliar spray in hybrid poplar plantations. CORAGEN® works best when insects are actively feeding on treated foliage. Application timing should be at first sign of economic damage.

Apply CORAGEN® at 2.5 to 5.0 fluid ounces per acre with properly calibrated ground or air equipment using sufficient water to obtain thorough coverage. For aerial application, use a spray volume of 5 gallons per acre or higher. A second application can be made as needed, but the second application must be at least 3 days after the first foliar application.

CORAGEN® can also be applied at 5.0 to 7.5 fluid ounces per acre via drip irrigation for suppression of Cottonwood Leaf Beetle. Application via drip irrigation works best during the first three years of plantation establishment. CORAGEN® must be applied uniformly in the root zone or poor performance will result. Consult the CORAGEN® section 3 label for more detailed instructions regarding application via drip irrigation systems. A second application can be made as needed, but the second application must be at least 10 days after the first drip application.

For resistance management, CORAGEN® is a group 28 insecticide. Do not make more than two successive applications of CORAGEN® before switching to an insecticide with a different mode-of-action.

RESTRICTIONS

Restricted-entry interval (REI) is 4 hours.

Do not apply more than 15.4 fluid ounces (0.20 lb a.i.) per calendar year by any combination of drip chemigation and foliar application.

IMPORTANT

Before using CORAGEN®, read and follow all applicable directions, restrictions, and precautions on the EPA-registered label.

This bulletin contains new or supplemental instructions for use of this product that does not appear on the package label. Follow the directions carefully.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.