



**DuPont Crop
Protection**

SUPPLEMENTAL LABELING

**DUPONT™ PREVATHON®
INSECT CONTROL
WITH ACTIVE INGREDIENT
RYNAXYPYR® FOR USE ON
TEFF AND QUINOA**

DUPONT™ PREVATHON® INSECT CONTROL

EPA Reg. No. 352-844

This supplemental label expires on 12/31/21 and must not be used or distributed after this date.

FOR FOLIAR INSECT CONTROL ON TEFF AND QUINOA

Not for sale, sale into, distribution and/or use in Nassau, Suffolk, Kings, and Queens counties of New York State

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This supplemental labeling must be in the possession of the user at the time of pesticide application.

PREVATHON® must be used only in accordance with the directions of this supplemental label and with the directions found on the container label. Always read this label and the entire label on the product container before using this product.

APPLICATION

Apply at the specified rates when insect populations reach locally determined economic thresholds. Consult the cooperative extension service, professional consultants or other qualified authorities to determine appropriate threshold levels for treatment in your area.

Apply follow-up treatments of PREVATHON®, as specified, to keep pest populations within threshold limits. Use sufficient water to obtain thorough, uniform coverage.

IMPORTANT

Before using PREVATHON®, read and follow all applicable directions, restrictions and precautions on the EPA Registered label. This bulletin contains new or supplemental instructions for use of this product, which do not appear on the EPA registered package label.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with such non-labeled use.

Crop	Application Method	Target Pest	DUPONT™ PREVATHON® RATE		Last Application (Days to Harvest)	REI (Hours)
			Lb ai per acre	fluid ounces product per acre		
Teff	FOLIAR	Corn earworm Beet armyworm European corn borer Fall armyworm Grasshoppers Sorghum webworm Southwestern corn borer Sugarcane borer True armyworm	0.047 - 0.067	14.0 - 20.0	14	4
	USE RESTRICTIONS Make no more than 4 applications per acre per crop. Minimum interval between treatments is 7 days. Do not apply more than 60 fl oz PREVATHON® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per year.					
Forage, Fodder, and Straw of Teff	FOLIAR	Corn earworm Beet armyworm European corn borer Fall armyworm Grasshoppers Sorghum webworm Southwestern corn borer Sugarcane borer True armyworm	0.047 - 0.067	14.0 - 20.0	14	4
	USE RESTRICTIONS Make no more than 4 applications per acre per crop. Minimum interval between treatments is 7 days. Do not apply more than 60 fl oz PREVATHON® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per year.					
Quinoa	FOLIAR	Corn earworm Beet armyworm European corn borer Fall armyworm Grasshoppers Sorghum webworm Southwestern corn borer Sugarcane borer True armyworm	0.047 - 0.067	14.0 - 20.0	14	4
	USE RESTRICTIONS Make no more than 4 applications per acre per crop. Minimum interval between treatments is 7 days. Do not apply more than 60 fl oz PREVATHON® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per year.					