

SUPPLEMENTAL LABELING

**DuPont Crop
Protection**

**DUPONT™ ALLY® XP
HERBICIDE**

**ROTATION INTERVAL TO SOYBEANS WITH
BOLT™ TECHNOLOGY**

DUPONT™ ALLY® XP HERBICIDE

EPA REG. NO. 352-435

ROTATION INTERVAL TO SOYBEANS WITH BOLT™ TECHNOLOGY

This supplemental label expires on March 17, 2019 and must not be used or distributed after this date.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CROP ROTATION

Minimum Rotation Intervals

Minimum rotation intervals* are determined by the rate of breakdown of ALLY® XP applied. ALLY® XP breakdown in the soil is affected by soil pH, presence of soil microorganisms, soil temperature, and soil moisture. Low soil pH, high soil temperature, and high soil moisture increase ALLY® XP breakdown in soil, while high soil pH, low soil temperature, and low soil moisture slow ALLY® XP breakdown.

Of these 3 factors, only soil pH remains relatively constant. Soil temperature, and to a greater extent, soil moisture, can vary significantly from year to year and from area to area. For this reason, soil temperatures and soil moisture should be monitored regularly when considering crop rotations.

*The minimum rotation interval represents the period of time from the last application to the anticipated date of the next planting.

Soil pH Limitations

Do not use ALLY® XP on soils having a pH above 7.9, as extended soil residual activity could extend crop rotation intervals beyond normal. Under certain conditions, ALLY® XP could remain in the soil for 34 months or more, injuring wheat and barley. In addition, other crops planted in high-pH soils can be extremely sensitive to low concentrations of ALLY® XP.

Checking Soil pH

Before using ALLY® XP, determine the soil pH of the areas of intended use. To obtain a representative pH value for the test area, take several 0" to 4" samples from different areas of the field and analyze them separately. Consult local extension publications for additional information on recommended soil sampling procedures.

ROTATION INTERVALS FOR CROPS IN NON-IRRIGATED LAND FOLLOWING USE OF ALLY® XP HERBICIDE

States	Crop	Soil pH	Application Rate (oz/A)	Minimum Rotation Interval (months)
Colorado Kansas Nebraska North Dakota (W. of Hwy. 1) Oklahoma South Dakota Texas	BOLT™ technology Soybeans	7.9 or lower	0.1	4

Unless indicated, trademarks with ®, TM or SM are trademarks of DuPont, Pioneer or affiliates. © 2016 DuPont.

© 2016 E. I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Road, Wilmington, Delaware 19805

IMPORTANT

BEFORE USING ALLY® XP HERBICIDE, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

Derived from SL-1996MSTR 031516 03-17-16