

SUPPLEMENTAL LABELING

**DuPont Crop
Protection**

**DUPONT™ GLEAN® XP
HERBICIDE**

**ROTATION INTERVAL TO SOYBEANS WITH
BOLT™ TECHNOLOGY**

DUPONT™ GLEAN® XP HERBICIDE

EPA REG. NO. 352-653

ROTATION INTERVAL TO SOYBEANS WITH BOLT™ TECHNOLOGY

This supplemental label expires on March 17, 2019 and must not be used or distributed after this date.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

CROP ROTATION

Minimum Rotation Intervals

Minimum rotation intervals* are determined by the rate of breakdown of GLEAN® XP applied. GLEAN® XP breakdown in the soil is affected by soil pH, soil temperature, and soil moisture. Low soil pH, high soil temperature, and high soil moisture increase GLEAN® XP breakdown in soil, while high soil pH, low soil temperature, and low soil moisture slow GLEAN® XP breakdown.

Of these 3 factors, only soil pH remains relatively constant. Soil temperature, and to a greater extent, soil moisture, can vary significantly from year to year and from area to area. For this reason, soil temperatures and soil moisture should be monitored regularly when considering recropping.

*The minimum rotation interval represents the period of time from the last application to the anticipated date of the next planting.

Soil pH Limitations

GLEAN® XP should not be used on soils having a pH above 7.9, as extended soil residual activity could extend crop rotation intervals beyond normal, and under certain conditions, could injure wheat, barley, or oat. In addition, other crops planted in high-pH soils can be extremely sensitive to low concentrations of GLEAN® XP.

Checking Soil pH

Before using GLEAN® XP, determine the soil pH of the areas of intended use. To obtain a representative pH value for the test area, take several 0" to 4" samples from different areas of the field and analyze them separately. Consult local extension publications for additional information on recommended soil sampling procedures.

NONCEREAL CROPS -- RECROPPING INTERVALS -- NON IRRIGATED LAND

States	Crop	Soil pH	Application Rate (oz/A)	Minimum Rotation Interval (months)
Arkansas Colorado Georgia Kansas Maryland Nebraska North Carolina Oklahoma South Carolina Texas Virginia	BOLT™ technology soybeans*	7.5 or lower	0.17 to 0.33	4

* Under certain conditions (such as drought, prolonged cold weather, pH variability in the fields) temporary discoloration and/or crop injury may occur to BOLT™ technology soybeans planted after DuPont™ GLEAN® XP applications. These intervals do not apply to crops grown for seed. These intervals may also be used for irrigated land.

© 2016 E. I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Road, Wilmington, Delaware 19805

IMPORTANT

BEFORE USING GLEAN® XP HERBICIDE, READ AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS ON THE EPA-REGISTERED LABEL.

This bulletin contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully.

This labeling must be in the possession of the user at the time of pesticide application.

Unless indicated, trademarks with ®, TM or SM are trademarks of DuPont or its affiliates. © 2016 DuPont.

Derived from SL-1997 031716 03-17-16