



**SPECIAL LOCAL NEED  
FIFRA Section 24(C)LABEL**

**DuPont Crop  
Protection**

**DUPONT™CORAGEN®  
INSECT CONTROL  
HYBRID POPLAR PLANTATIONS  
STATE OF WASHINGTON**

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

**DUPONT™ CORAGEN® INSECT CONTROL**

**EPA Reg. No. 352-729  
EPA SLN NO. WA-160005**

**FOR USE ON HYBRID POPLAR PLANTATIONS  
FOR CONTROL OF COTTONWOOD LEAF BEETLES**

**THIS LABEL FOR DUPONT™ CORAGEN® INSECT CONTROL EXPIRES AND MUST NOT BE  
DISTRIBUTED OR USED IN ACCORDANCE WITH THIS SLN REGISTRATION AFTER DECEMBER 31, 2021**

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

This labeling must be in the possession of the user at the time of application.

Follow all applicable directions, restrictions, Worker Protection Standard requirements and precautions on the EPA registered label.

**USE INFORMATION**

DuPont™ CORAGEN® Insect Control is a suspension concentrate that can be applied to control listed insect pests. CORAGEN® is diluted with water for application.

**APPLICATION INFORMATION, RATES AND TIMING**

CORAGEN® insect control is labeled for the control of Cottonwood Leaf Beetle (*Chrysomela scripta*) adults and larvae when applied as a foliar spray in hybrid poplar plantations. CORAGEN® works best when insects are actively feeding on treated foliage. Application timing should be at first sign of economic damage.

Apply CORAGEN® at 2.5 to 5.0 fluid ounces per acre with properly calibrated ground or air equipment using sufficient water to obtain thorough coverage. For aerial application, use a spray volume of 5 gallons per acre or higher. A second application can be made as needed, but the second application must be at least 3 days after the first foliar application.

CORAGEN® can also be applied at 5.0 to 7.5 fluid ounces per acre via drip irrigation for suppression of Cottonwood Leaf Beetle. Application via drip irrigation works best during the first three years of plantation establishment. CORAGEN® must be applied uniformly in the root zone or poor performance will result. Consult the CORAGEN® section 3 label for more detailed instructions regarding application via drip irrigation systems. A second application can be made as needed, but the second application must be at least 10 days after the first drip application.

For resistance management, CORAGEN® is a group 28 insecticide. Do not make more than two successive applications of CORAGEN® before switching to an insecticide with a different mode-of-action.

**RESTRICTIONS AND PRECAUTIONS**

- Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 4 hours.
- Do not apply more than 15.4 fluid ounces (0.20 lb a.i.) of CORAGEN® per calendar year by any combination of drip chemigation and foliar application.
- This use is only for hybrid poplars grown on land classified as cropland by the Farm Service Agency.

- This pesticide is toxic to aquatic invertebrates, oysters, and shrimp. CORAGEN® should not be used under this SLN label where impact on listed threatened or endangered species is likely. You may refer to the WSDA Endangered Species Program web site at [agr.wa.gov/PestFert/NatResources/EndangSpecies.aspx](http://agr.wa.gov/PestFert/NatResources/EndangSpecies.aspx), or contact the Washington Department of Fish & Wildlife, National Marine Fisheries Service (NOAA Fisheries) or US Fish & Wildlife Service for information regarding aquatic species listed as threatened or endangered. Consult the federal label for additional restrictions and precautions to protect aquatic organisms.

- Surface Water Advisory - This product may impact surface water quality due to runoff of rain water. This is especially true for poorly draining soils and soils with shallow ground water. This product is classified as having high potential for reaching surface water via runoff for several months or more after application. A level, well-maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential loading of chlorantraniliprole from runoff water and sediment. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours.
- Ground Water Advisory - This chemical has properties and characteristics associated with chemicals detected in ground water. This chemical may leach into ground water if used in areas where soils are permeable, particularly where the water table is shallow.

#### WSDA Chemigation Guidance:

- An inspection port or a direct access point is required, and it must be positioned immediately upstream of the irrigation mainline check valve and be of sufficient size to allow visual and manual inspection of the check valve and low pressure drain. The inspection port or access point must have a minimum diameter of four inches, unless an alternative access system is approved by WSDA (WAC 16-202-1012[1]).
- The chemigation application tank cannot be placed within 20 feet of the wellhead or other sensitive areas. Mixing or loading activities cannot occur within 20 feet of the wellhead or other sensitive areas (WAC 16-202-1008[1]).
- WSDA Chemigation Rules (WAC 16-202-1001 through WAC 16-202-1024), and information on USEPA Authorized Alternative Chemigation Safety Equipment, Distribution Uniformity and other chemigation topics are available on the WSDA website [agr.wa.gov/PestFert/ChemFert/default.aspx](http://agr.wa.gov/PestFert/ChemFert/default.aspx).

#### WSDA Container Disposal Guidance:

Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the WSDA Waste Pesticide Program web site at [agr.wa.gov/PestFert/Pesticides/WastePesticide.aspx](http://agr.wa.gov/PestFert/Pesticides/WastePesticide.aspx). Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington.

## IMPORTANT

Before using CORAGEN®, read and follow all applicable directions, restrictions, and precautions on the EPA-registered label.

This Special Local Need (SLN) Registration contains new or supplemental instructions for use of this product that does not appear on the package label. Follow the directions carefully.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

**For product information call 1-888-6-DUPONT**

**© 2016 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Road, Wilmington, DE 19805 U.S.A.**