

## 2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA



### DUPONT™ PREVATHON® INSECT CONTROL

EPA Reg. No. 352-844

### DUPONT™ STEWARD® EC INSECTICIDE

EPA Reg. No. 352-638

## FOR SUPPRESSION OF DECTES STEM BORER IN SOYBEAN IN THE STATE OF TEXAS

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Dupont™ PREVATHON® insect control is recommended for suppression of Dectes stem borer (*Dectes texanus*) in soybeans. Apply PREVATHON® at 20 fluid ounces per acre (0.067 lb ai/acre) as a foliar spray. As an alternative, apply PREVATHON® at 14 fluid ounces per acre (0.047 lb ai/acre) in a tank mix with STEWARD® EC at 6 fluid ounces per acre (0.059 lb ai/acre) as a foliar spray. Use a minimum of 10 gallons per acre (GPA) of water for ground applications or 3 GPA for aerial applications. For best results, use 20 GPA by ground or 5 GPA by air. See product labels for chemigation instructions.

To minimize crop damage by the pest, apply at the onset of adult beetle flight. Ensure thorough spray coverage and make application to soybeans prior to egg laying. For best results, regular scouting using a sweep net is necessary to identify the emergence and infestation of adult beetles. If regular scouting is not used, apply at 1500 Growing Degree Days (GDD) in Nebraska and northern Kansas or consult with your local agricultural advisor for advice on application timing. Continued scouting should be used to track the duration of the emergence period. A second application may be necessary at 3 to 4 weeks after the initial application if adults continue to emerge over an extended period.

### PRECAUTIONS

Do not apply more than 60 fluid ounces of PREVATHON® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per year. Do not apply more than 45 fluid ounces of STEWARD® EC or 0.44 lb a.i. of indoxacarb containing products per acre per year. Do not make more than 4 applications per acre per crop.

For PREVATHON®, the minimum interval between treatments is 3 days and preharvest interval is 1 day. If STEWARD® EC is used, the minimum interval between treatments is 5 days and preharvest interval is 21 days.

For PREVATHON® the field re-entry interval is 4 hours. If STEWARD® EC is used, the field re-entry interval is 12 hours.

Read and follow all applicable directions, restrictions and precautions.

#### IMPORTANT NOTICE

**BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY, AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.**

It is a violation of Federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This Bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This Bulletin must be in the possession of the User at the time of application. **Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using the DuPont product(s) covered by this Bulletin. DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with such uses not specified by the label or DuPont 2(ee) instructions. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability of the Section 3 Federal product label.**

Copies of DuPont product labels are available by calling 1-888-6-DUPONT  
or at <http://cropprotection.dupont.com/>

© 2017 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd., Wilmington, DE 19805 U.S.A