

**SPECIAL LOCAL NEED
24(C)LABELING**

**DuPont Crop
Protection**

**DUPONT™ UPBEET®
HERBICIDE
FOR USE ON GARDEN BEETS
IN THE STATE OF NEW YORK**

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF NEW YORK

DUPONT™ UPBEET® HERBICIDE

EPA Reg. No. 352-569

SLN No. NY-120015

**This label for DuPont UPBEET® Herbicide expires and must not be distributed or used
in accordance with this SLN registration after April 30, 2022**

FOR POSTEMERGENCE WEED CONTROL IN GARDEN BEETS

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Follow all applicable directions, restrictions and precautions including statements pertaining to the Worker Protection Standards on the EPA registered UpBeet® label.

This SLN label contains new or supplemental instructions for use of this product which do not appear on the EPA-registered package label. Follow the instructions carefully. This labeling must be in the possession of the user at the time of pesticide application.

Do not use this product until you have read the entire label. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

Do not use in Suffolk and Nassau Counties, Long Island, New York.

PRODUCT INFORMATION

DuPont™ UPBEET® herbicide may be used in Garden Beets for selective postemergence control of broadleaf weeds including wild mustard, shepard's-purse, and velvetleaf.

This product is a water dispersible granule containing 50% active ingredient by weight.

APPLICATION INFORMATION

Apply UPBEET® by ground only at a broadcast rate of 0.5 ounces per acre, starting when garden beets are at the 2 to 4 leaf stage. Additional applications may be made at the 4 to 6 leaf stage and at the 6 to 8 leaf stage.

The total amount of UPBEET® applied must not exceed 1.5 ounces per acre per growing season.

For best results apply UPBEET® to small actively growing weeds when the temperature is between 40F and 75F.

Do not treat when frost is expected in the hours following application. If high temperatures are expected during the day, treatment should be applied early in the morning or in the evening, so that a period of moderate temperatures (lower than 75F) will follow the application.

In some cases, discoloration and even slowed growth may be observed. These symptoms are usually temporary.

For product information call 1-888-6-DUPONT

© 2017 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Road, Wilmington, DE 19805

Refer to the Section 3 label for additional application instructions and use precautions.

Apply no later than 30 days before harvest.

WEED STAGE AT APPLICATION

Weeds should be actively growing and not under stress. Applications made to larger weeds or to weeds under stress may result in unsatisfactory control.

Since UPBEET® has little to no soil activity, only weeds that have emerged above the soil surface will be controlled.

Use sequential applications to control new weed flushes. The growth stage of weeds at application is very important for satisfactory control. For best results, apply to small, emerged weeds between the cotyledon and two true leaf stage.

TANK MIX OPTIONS

UPBEET® may be tank mixed with other suitable registered herbicides to broaden the weed control. DuPont recommends that you first consult your state experiment station, university, or extension agent, Agricultural dealer, or DuPont representative as to the potential for crop injury before using the mixture. If no information is available, limit the initial use of UPBEET® and the tank mix herbicide to a small area.

UPBEET® may also be mixed with other suitable registered fungicides or insecticides labelled for use on garden beets.

Read all label precautions for tank mix partners prior to use. Follow all manufacturers' label directions for the companion product. If these directions conflict with this label, do not tank mix with UPBEET®.

SPRAY ADDITIVES

Nonionic Surfactant (NIS) and Crop Oil Concentrate (COC)

- A tankmix with 0.25 % volume/volume (2 pt per 100 gal of spray solution) of non-ionic surfactant or crop oil concentrate may be added. Addition of surfactant may increase crop response. First time users should limit the use of surfactant to a small test area to determine if crop response is acceptable.
- Surfactant products must contain at least 60% nonionic surfactant with a hydrophilic/lipophilic balance (HLB) greater than 12.

IMPORTANT NOTICE

BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

It is a violation of federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This bulletin should be in the possession of the user at the time of application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.

Copies of DuPont product labels are available by calling 1-888-6-DUPONT
or at <http://cropprotection.dupont.com/>