

2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA



DUPONT™ AVAUNT® INSECTICIDE

EPA Reg. No. 352-597

FOR CONTROL OF KATYDID* NYMPHS IN STONE FRUIT AND GRAPE IN THE STATE OF CALIFORNIA

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product. For any requirements specific to your State or Tribe; consult the agency responsible for pesticide regulation. Do not apply this product in any way that will contact workers or other persons; either directly or through drift. Only protected handlers may be in the area during application.

DuPont™ AVAUNT® insecticide is recommended as a foliar application to control katydid nymphs in stone fruit and grape.

Apply AVAUNT® at 6.0 ounces/acre as a foliar spray at first indication of katydid nymphs. Allow 5 to 7 days to achieve maximum results.

Make repeat applications to stone fruit on a 7 to 10 day schedule if monitoring indicates continued feeding activity. Minimum spray interval for grape is 21 days.

Correct timing of spray applications is to the early nymphal stages; apply in sufficient water to obtain thorough and uniform coverage of foliage to achieve best results. Make application before pests reach damaging levels. If AVAUNT® is applied in two consecutive applications; it is recommended to switch to a different mode-of-action insecticide before making another application with AVAUNT® to reduce the risk of insect resistance. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action threshold levels for this pest. For best results, use an adjuvant to help increase coverage, penetration and thus performance.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with uses not specified by label or DuPont 2(ee) instructions.

PRECAUTIONS

Stone Fruit

Make no more than 3 applications prior to hand-thinning. No hand-thinning after the 4th application. Make no more than 4 applications per season.

* Forktailed bush katydid (*Scudderia furcata*) Angularwinged katydid (*Microcentrum retinerve*)

IMPORTANT NOTICE

BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY, AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

It is a violation of Federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This Bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This Bulletin must be in the possession of the User at the time of application. **Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using the DuPont product(s) covered by this Bulletin. DuPont will not be responsible for losses or damages resulting from use of this product in any manner not specifically stated by DuPont on the Section 3 label, or other labels, or this Bulletin. User assumes all risks associated with any such nonspecified use. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability of the Section 3 Federal product label.**

Copies of DuPont product labels are available by calling 1-888-6-DUPONT
or at <http://cropprotection.dupont.com/>

© 2017 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd., Wilmington, DE 19805 U.S.A

Do not apply more than 24 oz. AVAUNT® or 0.44 lbs ai of indoxacarb containing products per acre per year.
Do not apply dilute applications of more than 200 gal water per acre.
The minimum interval between sprays is 7 days.
Preharvest interval (PHI): Do not apply within 14 days of harvest.
The field re-entry interval (REI) is 12 hours.

Grapes

Make no more than 2 applications per season
Do not apply more than 12 oz. AVAUNT® or 0.22 lbs ai of indoxacarb containing products per acre per year.
The minimum interval between sprays is 21 days.
Preharvest interval (PHI): Do not apply within 7 days of harvest
The field re-entry interval (REI) is 12 hours.