

2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA



DUPONT™ AVAUNT® INSECTICIDE

EPA Reg. No. 352-597

FOR CONTROL OF LIGHT BROWN APPLE MOTH ON GRAPE, POME FRUITS AND STONE FRUITS IN THE STATES OF CALIFORNIA, OREGON AND WASHINGTON

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product. For any requirements specific to your State or Tribe; consult the agency responsible for pesticide regulation. Do not apply this product in any way that will contact workers or other persons; either directly or through drift. Only protected handlers may be in the area during application.

DuPont™ AVAUNT® insecticide is recommended for the control of Light Brown Apple Moth (*Epiphyas postvittana*) in Grapes, Pome Fruit and Stone Fruits.

Apply DuPont AVAUNT® Insecticide at 5.0 to 6.0 ounces per acre as a foliar spray using properly calibrated ground spray equipment.

Make the first application at initiation of egg hatch, at appearance of small larvae or at first signs of infestation for each generation. Use the higher application rate for moderate to heavy insect pressure. Make application before pests reach damaging levels. Monitor fields and make an additional application if populations rebuild to potentially damaging levels. Apply in sufficient water to obtain thorough and uniform cover of foliage. Consult your state cooperative extension service, professional consultants or other qualified authorities to determine appropriate action threshold levels for this pest.

For best results, use an adjuvant to help increase coverage, penetration and thus performance.

Refer to product label for DuPont AVAUNT® insecticide for additional general information and Precautions for each crop listed on this recommendation.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with uses not specified by the label or DuPont 2(ee) instructions.

Restrictions

Grapes:

Make no more than 2 applications per season

Do Not apply more than 12 oz AVAUNT® or 0.22 lb ai of indoxacarb containing products per acre per year.

Minimum interval between sprays is 21 days.

The field re-entry interval (REI) is 12 hours.

Preharvest interval (PHI): Do not apply within 7 days of harvest.

IMPORTANT NOTICE

BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY, AND FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

It is a violation of Federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This Bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This Bulletin must be in the possession of the User at the time of application.

Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using the DuPont product(s) covered by this Bulletin. DuPont will not be responsible for losses or damages resulting from use of this product in any manner not specifically stated by DuPont on the Section 3 label, or other labels, or this Bulletin. User assumes all risks associated with any such nonspecified use. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability of the Section 3 Federal product label.

Copies of DuPont product labels are available by calling 1-888-6-DUPONT
or at <http://cropprotection.dupont.com/>

© 2017 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd., Wilmington, DE 19805 U.S.A

Pome fruits (including Pear) and Stone fruits:

Make no more than 3 applications prior to hand-thinning. No hand-thinning after the 4th application.

Make no more than 4 applications per season.

Do Not apply more than 24 oz AVAUNT® or 0.44 lb ai of indoxacarb containing products per acre per year.

Do not apply dilute applications of more than 200 gal water per acre.

Minimum interval between treatments is 7 days.

The field re-entry interval (REI) is 12 hours.

Preharvest interval (PHI) for Pear, Pome and Stone Fruits:

Pome fruits (except Pears) and Stone fruits: Do not apply within 14 days of harvest.

Pears: Do not apply within 28 days of harvest.