

## 2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA



### **DUPONT™ CORAGEN® INSECT CONTROL** EPA Reg. No. 352-729

## **FOR CONTROL OF GRASSHOPPERS IN GRASS FORAGE FODDER AND HAY (RANGELAND & PASTURE GRASS) IN THE STATE OF TEXAS**

**This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.**

### **DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Always read the FIFRA 2(ee) bulletin and the entire label on the product container before using this product. For any requirements specific to your State or Tribe; consult the agency responsible for pesticide regulation. Do not apply this product in any way that will contact workers or other persons; either directly or through drift. Only protected handlers may be in the area during application.

DuPont™ CORAGEN® insecticide is recommended for the control of grasshoppers in grass forage, fodder, and hay: any grass, Gramineae family (rangeland & pasture grass).

Apply CORAGEN® at 2.0 to 4.0 fluid ounces per acre (0.026 - 0.052 LB A.I. PER ACRE) as a foliar spray via aerial application or ground application. For best results with foliar sprays, add Methylated Seed Oil (MSO) adjuvant at 1 gallon per 100 gallons of spray volume (1% v/v).

Apply when grasshopper populations reach local established thresholds to prevent crop damage. Correct timing of spray applications to nymphal stages and thorough coverage is critical to achieve optimum control. For best results applications should be made when eggs have hatched and the majority of the grasshopper population is 2nd – 3rd instar nymphs. Read and follow all spray drift management guidance in the Federal Section 3 CORAGEN® label.

Once grasshoppers contact and/or ingest CORAGEN® there will be rapid feeding cessation; full expression of mortality may take a week or more.

Do not make more than two sequential applications of CORAGEN® before rotating to another registered insecticide having a different mode-of-action.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with uses not specified by the label or DuPont 2(ee) instructions.

### **PRECAUTIONS**

Do not apply more than 15.4 fluid ounces (0.2 lb a.i.) of CORAGEN® per acre per year.

Do not make more than 2 applications per crop. The minimum retreatment interval is 7 days.

The preharvest interval is 0 days. The field re-entry interval is 4 hours.

Read and follow all applicable directions, restrictions, and precautions on the EPA-registered label.

(Replaces R-1481 052714 05-28-14)

#### **IMPORTANT NOTICE**

**BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.**

It is a violation of federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This bulletin should be in the possession of the user at the time of application.

**Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.**

Copies of DuPont product labels are available by calling 1-888-6-DUPONT  
or at <http://cropprotection.dupont.com/>

© 2017 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd., Wilmington, DE 19805 U.S.A

R-1858 051717 05-23-17

Page 1 of 1

Expires 12/31/18