

## 2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA



### DUPONT™ CORAGEN® INSECT CONTROL

EPA Reg. No. 352-729

### FOR CONTROL OF CERTAIN PESTS IN GRASS GROWN FOR SEED IN THE STATES OF IDAHO, OREGON, AND WASHINGTON

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

#### DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product. For any requirements specific to your State or Tribe; consult the agency responsible for pesticide regulation. Do not apply this product in any way that will contact workers or other persons; either directly or through drift. Only protected handlers may be in the area during application.

DuPont Coragen® insect control is recommended at 3.5 to 5.0 fluid ounces per acre (0.045 - 0.065 lb a.i. per acre) for the control of Armyworms, Cutworms, and Sod Webworms. For suppression of Billbug grubs and European Crane Fly larvae, apply Coragen® at 5.0 fluid ounces/acre (0.065 lb a.i. per acre).

#### Method of Application

For control of Armyworms and Cutworms, apply Coragen® as a thorough coverage foliar spray using properly calibrated ground equipment in a minimum of 10 gallons per acre, or via overhead chemigation in 0.10 to 0.20 acre inch of water.

For foliar sprays, increase the spray volume to compensate for the amount of foliage present. For maximum spray penetration in to the root crown area, the use of a silicone surfactant may be useful. For best results with foliar spray applications, delay the next irrigation for at least 24 hours.

For control of Sod Webworm larvae, and suppression of Billbug grubs and Crane Fly larvae, it is important to move the Coragen® into the grass root zone. This is best achieved by applying via overhead chemigation in 0.25 to 0.50 acre inch of water, or by immediately following a foliar spray application with 0.25 to 0.50 acre inch of water.

#### Application Timing

For control of Armyworms, Cutworms, and Sod Webworms, apply at first sign of economic crop damage.

For suppression of European Crane Fly larvae apply between September and early November.

For suppression of Billbug grubs, apply when overwintered adult Billbugs are first observed. This will usually occur in late April or early May.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with uses not specified by the label or DuPont 2(ee) instructions.

#### IMPORTANT NOTICE

**BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.**

It is a violation of federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This bulletin should be in the possession of the user at the time of application.

**Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.**

Copies of DuPont product labels are available by calling 1-888-6-DUPONT  
or at <http://cropprotection.dupont.com/>

© 2017 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd., Wilmington, DE 19805 U.S.A

**Precautions**

Read and follow all instructions and precautions on the federal section 3 label before using Coragen® on grass grown for seed.

The preharvest interval is zero days.

The minimum re-entry interval is 4 hours.

Do not apply more than 5 fluid ounces per acre per treatment.

Do not make more than 4 applications per crop.

The minimum interval between applications is 7 days.

Do not apply more than 15.4 fluid ounces per acre (0.2 lb a.i. per acre) per year.