

2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA



DUPONT™ CORAGEN® INSECT CONTROL

EPA Reg. No. 352-729

FOR USE VIA DRIP CHEMIGATION APPLICATION IN GLOBE ARTICHOKE FOR CONTROL OF BEET ARMYWORM AND CABBAGE LOOPER IN THE STATE OF CALIFORNIA

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. Always read this FIFRA 2(ee) bulletin and the entire label on the product container before using this product. For any requirements specific to your State or Tribe; consult the agency responsible for pesticide regulation. Do not apply this product in any way that will contact workers or other persons; either directly or through drift. Only protected handlers may be in the area during application.

DuPont™ Coragen® insect control must be used only in accordance with directions on this bulletin or in separately published DuPont labeling.

DuPont™ Coragen® insect control is recommended for the control of beet armyworm and cabbage looper via drip chemigation in globe artichoke.

Read the chemigation section of the federal label for more information on how to use via drip chemigation.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with uses not specified by the label or DuPont 2(ee) instructions.

Crop	Application Method	Target Pest	DUPONT™ CORAGEN® RATE		Last Application (days to Harvest)	REI (Hours)
			Lb. ai per acre	Fluid ounces product per acre		
Artichoke, globe	Drip chemigation	Beet armyworm Cabbage looper	0.045-0.098	3.5 - 7.5	3	4
<p>USE RESTRICTIONS Begin drip application when populations first appear. CORAGEN® must be applied uniformly in the root zone or poor performance will result.</p> <p>Drip tape must be placed directly underneath a single row to ensure CORAGEN® is applied in the root zone.</p> <p>During the first half of the crop growing cycle, allow 1-3 days for CORAGEN® to be translocated into the aerial portions and to fully protect. For applications made in the second half of the crop growing cycle: translocation of CORAGEN® into aerial portions of the plant may take up to 7-10 days.</p> <p>Do not make more than 2 drip chemigation applications of CORAGEN® per crop.</p> <p>Minimum application interval between drip chemigation treatments is 14 days.</p> <p>Apply no more than 15.4 fl oz CORAGEN® or 0.2 lb a.i. of chlorantraniliprole containing products per acre per calendar year.</p>						

IMPORTANT NOTICE

BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.

It is a violation of federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This bulletin should be in the possession of the user at the time of application. **Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with such uses not specified by the label or DuPont 2(e) instructions. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.**

Copies of DuPont product labels are available by calling 1-888-6-DUPONT
 or at <http://cropprotection.dupont.com/>