



## 2(ee) Recommendation

Use information in accordance with Section 2(ee) of FIFRA

### **DUPONT™ VERIMARK® INSECT CONTROL**

EPA Reg. No. 352-860

#### **FOR THE SUPPRESSION OF TOMATO SPOTTED WILT VIRUS INCIDENCE IN TOBACCO VECTORED BY THRIPS IN THE STATES OF GA, KY, NC, PA, SC, TN AND VA**

This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or approved by EPA.

#### **DIRECTIONS FOR USE**

It is a violation of federal law to use this product in a manner inconsistent with its labeling. For any requirements specific to your State or Tribe; consult the agency responsible for pesticide regulation. Do not apply this product in any way that will contact workers or other persons; either directly or through drift. Only protected handlers may be in the area during application.

DuPont™ Verimark® insect control must be used only in accordance with directions on this bulletin or in separately published DuPont labeling.

DuPont™ Verimark® insect control used in programs with imidacloprid is recommended for the suppression of tomato spotted wilt virus in tobacco when vectored by thrips.

Apply DuPont™ Verimark® insect control at 13.5 fluid ounces per acre as a transplant tray drench or transplant water application. An application of a registered product that contains imidacloprid as a tray drench or transplant water at labeled rates, typically 0.028 - 0.04 lb ai/1000 plants, should also be made. Verimark® and imidacloprid can be applied together using the same application method or separately using different application methods. When applying Verimark® as a tray drench, apply no earlier than 72 hours prior to planting in the field. Read and follow label directions for transplant tray drench and transplant water application methods as described on the section 3 label of Verimark® and the imidacloprid containing product.

DuPont™ Verimark® insect control impairs insect muscle function, resulting in rapid cessation of feeding and limiting the ability of insects to move, which can reduce transmission and spreading of insect vectored diseases.

Read and follow the protection of pollinators section of the Section 3 Verimark® label.

Read and follow all applicable directions, restrictions and precautions on the EPA-registered label for each product used.

DuPont will not be responsible for losses or damages resulting from the use of this product in any manner not specifically instructed by DuPont. User assumes all risks associated with uses not specified by the label or DuPont 2(ee) instructions.

#### **RESTRICTIONS**

Do not apply a total of more than 0.4 lb ai/A of CYAZYPYR® or cyantraniliprole containing products per acre per calendar year.

Re-entry interval is 12 hours.

Pre-harvest interval is 3 days.

**IMPORTANT NOTICE**

**BEFORE BUYING OR USING DUPONT PRODUCT(S) COVERED BY THIS BULLETIN, READ THIS BULLETIN, AND THE LABELS FOR ALL PRODUCTS REFERENCED IN THIS BULLETIN, IN THEIR ENTIRETY. FOLLOW ALL APPLICABLE DIRECTIONS, RESTRICTIONS AND PRECAUTIONS.**

It is a violation of federal law to use this or any other pesticide product in a manner inconsistent with its labeling. This bulletin contains supplemental instructions for use of the referenced DuPont product. Follow these instructions carefully. This bulletin should be in the possession of the user at the time of application.

**Read the Limitation of Warranty and Liability on the Section 3 Federal product label before buying or using THIS product. If terms are not acceptable, return the unopened package at once to Seller for full refund of purchase price paid. Otherwise, use by Buyer or any other User constitutes acceptance of the terms of the Limitation of Warranty and Liability on the Section 3 Federal product label.**

Copies of DuPont product labels are available by calling 1-888-6-DUPONT  
or at <http://cropprotection.dupont.com/>

© 2017 E.I. du Pont de Nemours and Company, Chestnut Run Plaza, 974 Centre Rd., Wilmington, DE 19805 U.S.A